80	Pacalatrician Management Group (RF) NPC	Vacci	na	tic	n	Sc	he	d	ule	2	02	41	More the O YEA of Excelled	ARS ence	
	Disease	Vaccine	Birth	6 weeks	10 weeks	14 weeks	6 months	9 months	12 months	15 months	18 months	6 years	9 years	12 years	
	ТВ	BCG	00											Eng.	
	Polio	bOPV	• •	0 0			(
	Diphtheria, Tetanus, Polio, Pertussis	Hexaxim		•	•						•			op	
	Haemophilus Influenzae, Hepatitis B	Infanrix hexa		8 weeks	12 weeks	16 weeks					•			P.	
	Pneumococcal OR OR	00140			•			•		15 months Official recomm	endations should l	oe taken into ac	count when immunisin	ng with Synflorix	
	Rotavirus OR	Rotarix toocine RotaTeq (Rotavirus Vaccine, Live, Oral, Pentavalent)			•										
	Measles Mumps Rubella ^{OR}	OMZTIA							MMR at 12 months			•	Give MMR and Varilrix or Onvara separately at		
	Chickenpox OR	Varil rix Description of the state of the st								:		•	6 years		
	Measles, Rubella	SII MR													
	Meningococcal	Menactra						•		•	For individual	s 2 - 55 years	administer 1 dose		
	Hepatitis A	Hayrix Avaxim 80							Н	lep A with MN	AR				
	Tetanus, Diphtheria Pertussis, Polio	boostrix tetra diphthoria, fotanus. acellular pertuasis (bancoroci) and inactivated polinivas.	Tetraxim at 6 years or Boostrix Tetra at 6 years Boostrix Tetra can be given at 12 years if not given at 6 years. Tdap IPV for Tdap (from 4 yrs. of age onwards)												
	「etanus, Diphtheria, Pertussis	TdaP hoostriv										•		•	
	Human Papilloma Virus OR	25	Cervarix from 9 – 14 years as 2 doses 6 months apart. State programme: Cervarix to Grade 4 girls in public schools 9 -14 years 2 doses 6 months apart. Gardasil from 9 – 13 years as 2 doses 6 moths apart or Gardasil 9 as 2 doses 5 -13 months apart												
	Influenza	Influvac	From 6 month	From 6 months, important for children in creche, chronic illness and respiratory problems. Start before the INFLUENZA season in MAY.											
	Respiratory Syncitial Virus	SYNAGIS' PALIVIZUMAB	RSV prophylaxis in high-risk infants-prevention of serious LRTI caused by RSV. Start January and end in May.												
	Pneumococcal	PNEUMOVAX 23 (pneumococcal vaccine, polyvalent, MSD)		Must have at least one dose of Conjugate PCV before Pneumovax 23. Pneumovax only in children older than 2 years with Immune Compromise or high risk of Pneumococcal infection. Two doses with 2nd dose 3-5 years after first.											
	Hepatitis B	Engerix-B	This vaccine is given to babies who are exposed to HbsAg at birth											150	
0	0	O CTAT	E EDIVACO	CINIES T	Reserve to the same		wilela C		-		a lance				

STATE EPI VACCINES These vaccines are available free from Government supplied clinics

RECOMMENDED OPTIONAL VACCINES Some of the vaccines in this schedule are only available from private clinics